

Prostate Cancer: Risk Assessment and Diagnostic Approaches

Leonard G. Gomella, MD,* Fernand Labrie, MD,[†] Eduard J. Gamito,[‡] Michael K. Brawer, MD[§]

*Thomas Jefferson University School of Medicine, Philadelphia, PA; [†]Laval University Medical Center, Quebec, Canada;

[‡]ANNs in CaP Project, Institute for Clinical Research, Washington, DC; [§]Northwest Prostate Institute, Seattle, WA

The successful treatment of prostate cancer relies on detection of the disease at its earliest stages. Although prostate-specific antigen (PSA)-based screening has been a significant advance in the early diagnosis of prostate cancer, identifying specific genetic alterations in a given family or patient will allow more appropriate screening for early disease. Mapping and identification of specific prostate cancer susceptibility genes is slowly becoming a reality. Other prostate cancer risks include a family history, race, and possibly serum markers such as insulin-like growth factor-I (IGF-I). Once a high-risk man is identified, transrectal ultrasound (TRUS)-guided biopsies are the standard to diagnose prostate cancer. Although TRUS is an advance over traditional digitally directed biopsies, it represents a random sampling of the prostate since most lesions cannot be visualized. Newer modalities such as ultrasound contrast agents, pattern recognition, and artificial neural networks (ANNs), applied to TRUS images, may improve diagnostic accuracy. If a man at risk for prostate cancer has undergone a negative TRUS biopsy, the decision for the need for additional biopsies is problematic. Use of PSA derivatives such as free and total PSA and the initial biopsy abnormalities such as atypia or high-grade prostatic intraepithelial neoplasia may define those patients in need of follow-up biopsy. [Rev Urol. 2001;3(suppl 2):S31-S38]

© 2001 MedReviews, LLC

Key words: IGF-I • Prostate cancer • Prostate biopsy • Susceptibility genes • Transrectal ultrasound • Ultrasound contrast agents

Improvements in screening and diagnosing prostate cancer have occurred with the use of prostate-specific antigen (PSA) and the development of gray-scale transrectal ultrasound (TRUS) biopsy techniques. As the human genome project moves forward, we may be able to identify specific abnormalities that may predispose to cancer and thus alter our screening approach. Likewise, new markers, exemplified by serum insulin-like growth factor-I (IGF-I), may predict future development of the disease. Enhancements in TRUS biopsy, including contrast agents, artificial neural networks, and the parameters for follow-up biopsy will improve the accuracy of our current diagnostic techniques. This report presents highlights of papers presented at

Table 1
Potential Genetic Alterations Under Study in Prostate Cancer

- | | |
|----------------------------|---|
| • <i>HPC1</i> at 1q24 | • <i>ELAC2</i> at 17p12 |
| • <i>PCAP</i> at 1q42 | • <i>BRCA1/BRCA2</i> |
| • <i>HPCX</i> at Xq27 | • 5 α -reductase type II gene (<i>SRD5A2</i>) at 2p23-22 |
| • <i>HPC20</i> at 20q13.18 | • Androgen receptor alterations (CAG, CCG repeats) |
| • <i>CAPB</i> at 1p36 | |

the 11th International Prostate Cancer Update held in Vail, Colorado that addressed these current issues in the early diagnosis of prostate cancer.

Prostate Cancer
Susceptibility Genes

The prevalence of prostate cancer varies markedly between different ethnic groups, with the highest frequency in African Americans and the lowest frequency in Asian populations. This ethnic disparity may be attributable to environmental factors, genetic factors, or both. Studies have investigated the role of family history as a risk factor for prostate cancer.^{1,2} Reports of families with large numbers of cases, as in the Utah Mormon population, have been key for exploring inherited prostate cancer.^{3,4} The risk of prostate cancer increases with younger age of onset in a relative or when the number of affected individuals in a familial cluster increases. This increased risk is strong evidence for a genetic component. For example, the brother of a man diagnosed at age 50 has a relative risk of 1.9 times of developing a prostate cancer compared with a brother of a case diagnosed at age 70.⁵ Having two or three first-degree relatives affected increases the relative risk by 4.9 and 10.9 times, respectively.⁶ Monozygotic twins have a fourfold increased rate of prostate cancer compared with dizygotic

twins.⁷ Estimates using the combined data from 44,788 sets of twins in Swedish, Danish, and Finnish registries suggest that up to 42% of all prostate cancer risk may be explained by inheritable factors.⁵ Identification of these specific genetic alterations that place a man at a higher risk of developing and dying of prostate cancer is an area of intense investigation that is further facilitated by the Human Genome Project. A list of potential genetic alterations in prostate cancer can be found in Table 1.

A correlation between CAG repeat length in the androgen receptor and

have been described and tested on independent data sets.¹¹⁻¹⁴ Another recent study presents significant evidence for linkage to a new locus, *HPC20* at 20q13.¹⁵ Of these, only the *HPC1* linkage has a reasonable level of independent confirmation; other studies found no significant evidence for linkage.^{3,16,17} Although the initial report of linkage to *HPC1* suggested that up to 34% of prostate cancer families could be related to this locus, a subsequent pooled analysis of 772 families demonstrated the proportion to be about 6%.¹⁸

Positional cloning techniques have identified *ELAC2*, a candidate prostate cancer susceptibility gene at 17p12.¹⁹ This is the first prostate cancer susceptibility gene cloned after a genome-wide scan of high-risk families. There is evidence that both frameshift and missense mutations are disease associated at this locus.

Familial co-segregation of breast and prostate cancer has also been reported. In breast/ovarian cancer families, male carriers of a deleterious

Estimates using the combined data from 44,788 sets of twins in Swedish, Danish, and Finnish registries suggest that up to 42% of all prostate cancer risk may be explained by inheritable factors.

age of onset of prostate cancer has been observed.⁸ A study of 587 individuals demonstrated an inverse relationship between CAG repeat length in the androgen receptor (AR) gene and risk of aggressive prostate cancer.⁹ Other studies support an association between reduced AR CAG repeat length and increased risk of prostate cancer, the length of the polymorphism GGC repeat in the AR being an additional potential risk factor.¹⁰

Four prostate cancer susceptibility loci (*HPC1* at 1q24, *PCAP* at 1q42, *HPCX* at Xq27, and *CAPB* at 1p36)

mutation in *BRCA1* or *BRCA2* have been shown to be at a three- and sevenfold risk of prostate cancer, respectively.^{20,21}

Prostate cancer is the cancer most sensitive to hormonal manipulation. Analyses of genes encoding proteins involved in androgen biosynthesis and action led to the observation of a significant association between common genetic variants and a susceptibility to prostate cancer. One such gene is the 5 α -reductase type II gene (*SRD5A2*), which catalyzes the conversion of testosterone into dihy-

Table 2
Published Studies on the Role of
Elevated IGF-I and Prostate Cancer Risk

| Study | Total no. in Study (Prostate Cancer/Control) | |
|---|---|-----------|
| Studies supporting an association | | |
| Chan et al, 1998 ²⁹ | 304 | (152/152) |
| Montzaros et al, 1997 ²⁸ | 104 | (52/52) |
| Wolk et al, 1998 ²⁷ | 434 | (210/224) |
| Studies not supporting an association | | |
| Kanety et al, 1992 ³⁰ | 34 | (24/10) |
| Cohen et al, 1992 ³¹ | 48 | (32/16) |
| Ho et al, 1997 ³² | 31 | (16/15) |
| Schaefer and Friedman, 1998 ³³ | 215 | (45/170) |
| Baffa et al, 2000 ³⁶ | 96 | (57/39) |
| Shariat et al, 2000 ³⁵ | 150 | (120/30) |

drotestosterone (DHT) and maps to 2p23-22.²²

The growing body of knowledge of the potential familial aspects of prostate cancer provides an important tool for understanding the mechanisms underlying the molecular basis of prostate cancer. Study of the genetic aspects of tumor initiation and progression will probably yield more effective screening, treatment, and prevention strategies in the future.

IGF-I as A Marker for Prostate Cancer

The IGF system consists of a series of ligands (IGF-I, IGF-2, insulin), binding proteins (IGF binding proteins 1-6), and receptors. IGF-IR (also known as the type I IGF receptor) binds both IGF-I and IGF-II and is a cell surface receptor in the tyrosine kinase class. The IGF-II receptor preferentially binds IGF-II but does not appear to transduce a mitogenic signal. IGF-I has been proved to be an important mitogenic and antiapoptotic peptide in many tumors, including breast, lung, and colorectal, as well as prostate

cancer.^{23,24} In vitro studies have established that human prostate cancer cell lines have functional IGF-I, with blockage of the receptor leading to cessation of growth.^{25,26} The presence of IGF-I in prostate cells and its potential role in the growth and development of cancer suggest that IGF-I may serve as a predictor of prostate cancer or potential target for prostate cancer therapy.

The prostate cancer literature presents conflicting evidence to the predictive value of IGF-I and the development of prostate cancer. Some studies show a positive relationship between IGF-I and prostate cancer, whereas others show an inverse relationship or no relationship at all (Table 2).

An association between elevated levels of plasma IGF-I and an increased risk of prostate cancer has been shown in a number of studies. There appears to be no correlation between IGF-I and benign prostatic hyperplasia (BPH), but increased levels of IGF-I (> 60 ng/mL), are suggested to be associated with an increased risk of prostate cancer. Three studies (by Wolk et al,²⁷ Montzaros et al,²⁸

and Chan et al²⁹) support a statistically significant association between elevated IGF-I levels and the risk of developing prostate cancer.

Other studies do not come to the same conclusion. These studies have compared prostate cancer patients with counterparts who have normal prostates and benign prostatic hypertrophy.³⁰⁻³⁶ Several of these studies have actually suggested an inverse relation between IGF-I levels and risk.^{33,36} In patients undergoing radical prostatectomy for localized disease, plasma IGF-I levels predicted neither organ-confined disease nor the risk of PSA progression and did not correlate with preoperative PSA level or final Gleason score.

There may be many reasons for these diverse observations: the specific IGF-I assay system used, treatment with hormonal therapy, and, most importantly, consideration of the effects of IGF binding proteins on serum IGF-I levels. The role of IGF-I as a risk factor for prostate cancer deserves continued study.

With inhibition of the IGF-IR pathway, normal cells appear to simply stop growing, but cancer cells appear to die rapidly through an apoptotic mechanism. Impairment of the IGF-IR function has dramatic effects on cancer cells growing in anchorage independence compared with cells growing in monolayer.³⁷ If IGF-I proves not to be a reliable marker for the disease, it may still have a role in therapy.

Contrast-Enhanced Prostate Biopsy

Years of experience have shown that TRUS-directed prostate biopsy, although very useful, has several limitations.^{38,39} Prospective TRUS imaging data have demonstrated that conventional gray scale is slightly superior to random chance in prostate cancer detection.⁴⁰ The trend is to increase the number of biopsies

in order to compensate for the limitations of the imaging alone.⁴¹ The traditional lesion-directed biopsy led to the development of the six-core, or sextant, biopsy technique. Today, at least eight to ten biopsies are recommended to sample the prostate gland more adequately. These additional biopsies tend to be laterally directed. Imaging of the prostate must be improved in order to enhance the detection of cancer.

Enhanced ultrasound techniques, such as color flow Doppler imaging (CDI) studies, have been investigated to improve the diagnostic ability of TRUS. Although studies suggest that CDI has potential prognostic significance, CDI can overlap with prostatitis and has low sensitivity in detection of tumor blood flow within prostate cancer. Alterations in the vascularity of prostate cancer have focused attention on ways to visualize characteristic vascular abnormalities, such as microbubble contrast agents. The use of intravenous microbubbles (average diameter 2 to 5 μ m leads to marked enhancement of the signal-to-noise ratio and enhanced visuali-

Today, at least eight to ten biopsies are recommended to sample the prostate gland more adequately.

zation on TRUS.⁴² Several vascular ultrasound agents are being studied for their use in prostate imaging; most such agents have demonstrated enhancement on gray-scale imaging (Table 3).

A study using EchoGen (Sonus Pharmaceuticals, Bothell, WA) consisted of 15 patients with a rising PSA; 14 of the 15 had a negative prior biopsy.⁴³ CDI TRUS was performed before and after EchoGen administration and correlated with sextant biopsies. Abnormal microvessel patterns were noted in eight

| Table 3 Some Vascular Ultrasound Contrast Agents Under Study in Imaging Prostate Cancer |
|--|
| <ul style="list-style-type: none">• Echovist (Schering, Berlin, Germany)• Levovist (Schering)• EchoGen (Sonus Pharmaceuticals, Bothell, WA)• Imagent US (Alliance Pharmaceutical, San Diego, CA)• Definity (DuPont Pharmaceuticals, Billerica, MA) |
| None of the agents are currently approved by the U.S. Food and Drug Administration for use in the evaluation of prostate cancer. |

patients; in two of these the lesions were malignant, one patient was diagnosed with prostatitis, and in two the lesions were benign. False-negative results were observed in three patients. Levovist (Schering AG, Berlin, Germany) was used in nine cases of prostatic cancer, with blood flow images enhanced in all cases.⁴⁴ In both studies, the authors concluded that contrast CDI is a promising technique that may allow for better imaging of blood flow and more accurate detection of early

contrast agents is dependent on the ability of these agents to traverse the tumor neovascularity safely without being destroyed. Unfortunately, conventional ultrasound systems deliver power levels sufficient to destroy microbubbles. A potential solution to this problem is the use of intermittent imaging. Standard gray-scale ultrasound image is refreshed at 30 frames per second; thus the amount of contrast agent available for each frame is that which enters the imaging plane in 1/30th of a second. This short period is usually not sufficient for contrast agents to enter small-diameter vessels. With intermittent imaging, the ultrasound beam is turned off for longer periods between each frame, allowing more contrast material to enter the imaging plane during this interscan period. An initial experience at Thomas Jefferson University with 26 subjects indicated that intermittent imaging might enhance visualization of malignant neovascularity, with several patients demonstrating enhancement of tumor foci not detected by conventional gray-scale imaging or CDI.

The group from Thomas Jefferson University has reported using Imagent (Alliance Pharmaceutical Corp., San Diego, CA) as a prostate contrast agent.⁴⁶ Twenty-six subjects

malignant lesions. Bogers et al⁴⁵ were the first to report the use of contrast-enhanced three-dimensional power Doppler angiography in the human prostate. With Levovist as a contrast agent, 18 patients with a suspicion of prostate cancer were evaluated; the study findings indicated that contrast-enhanced power Doppler and three-dimensional image reconstruction offers a useful imaging tool with good potential to improve prostate cancer detection in the future.

Adequate enhancement of ultrasound imaging using microbubble

with an elevated PSA and/or abnormal digital rectal examination (DRE) were studied. Continuous gray-scale, intermittent gray-scale, phase inversion gray-scale, and power Doppler sonography of the prostate were performed and correlated with sextant biopsy results. The study demonstrated significant visible enhancement ($P < .05$) after administration of Imagent.

In another study, 60 subjects were evaluated with conventional gray-scale, harmonic gray-scale, and power Doppler sonography.⁴⁷ The evaluation was repeated using IV Definity (DuPont Pharmaceuticals, Billerica, MA). Gray-scale imaging was performed in continuous mode and with intermittent imaging using interscan delay times of 0.5, 1.0, 2.0, and 5.0 seconds. Sextant biopsy sites were scored prospectively as benign or malignant on baseline imaging, and again during contrast-enhanced TRUS. Prostate cancer was present in 37 biopsy sites from 20 subjects. Baseline imaging demonstrated prostate cancer in 14 sites from 11 subjects. Contrast-enhanced TRUS demonstrated prostate cancer in 24 sites from 15 subjects. Each of the five subjects whose prostate cancer was missed had only a single positive biopsy score (Gleason score ≤ 6). The improvement in sensitivity from 38% at baseline to 65% with contrast was significant ($P < .004$). Using the contrast agent Definity along with TRUS improves sensitivity for detection of malignant foci within the prostate without substantial loss of specificity.

The currently available data suggest that ultrasound contrast agents may enhance our ability to identify specific foci of prostate cancer on TRUS. In particular, higher grade cancers may be more easily detected than low-grade lesions. These contrast agents may become a standard part of TRUS biopsy in the future.

Analysis of TRUS Images by Artificial Neural Networks

Another potential means of enhancing TRUS images and identifying malignant foci is the use of artificial neural networks. Automated image analysis, including pattern recognition and artificial neural networks (ANNs) applied to TRUS images, may successfully identify lesions that cannot be seen by the human eye. At present, such automated image analysis and pattern recognition are unavailable for existing TRUS systems.

ANNs, a type of artificial intelligence, are a software construct based roughly on the neural structure of the brain. Basic processing units called nodes simulate neurons, and weighted interconnections between the nodes simulate dendrites and

previously. This "trained" ANN will function in a way similar to a mathematical function with inputs analogous to independent variables and outputs analogous to dependent variables. The ANN's performance can be measured by calculating the sensitivity, specificity, negative predictive value (NPV), and positive predictive value (PPV) for a specific ANN output cutoff. The overall performance (ie, over all output cutoffs) may be quantified by generating a receiver operator characteristic (ROC) curve.

In 1992, the first efforts to analyze TRUS images of the prostate were published.⁴⁹ ANN was able to distinguish between prostatic and nonprostatic tissues in TRUS images. Investigators have used ANNs for spectral analysis of ultrasound RF

The currently available data suggest that ultrasound contrast agents may enhance our ability to identify specific foci of prostate cancer on TRUS.

axons.⁴⁸ The interconnection weights function as multipliers that simulate the connection strengths in the analogous biological model. The ANN is not programmed but learns by experience, via a "supervised learning" phase called training. Other types of ANNs may rely on an unsupervised learning method.⁴⁸ Cases that include inputs and known outputs, such as sets of clinical variables and a known pathological outcome, are presented to the ANN sequentially and repeatedly. A training algorithm automatically adjusts the connection weights, consequently changing the output values, to reduce errors between the actual ANN outputs and the expected outputs. As the ANN is trained, a set of connections is developed that allow for the largest number of correct predictions for the given training data set. The ANN is next validated with new cases not used

signals (pre-image) and ANNs to analyze ultrasound images of breast, colon, prostate, and other tissues, with promising results.^{50,51} Workers at the University of Kiel, Germany, have assembled a prospective library of prostate tissue types by gathering TRUS images prior to radical prostatectomy (RP) and comparing these with whole-mount pathology slides.⁵² In this work, an ANN was used to identify areas suspicious for cancer in a validation set of TRUS images. Preliminary data demonstrated that 99% of confirmed benign samples were correctly identified, with 79% of malignant lesions correctly classified. Ninety-seven percent of isoechoic cancers on TRUS were correctly classified by the ANN. Workers with the Artificial Neural Networks in Carcinoma of the Prostate (ANNs in CaP) Project (Crawford, Gamito, and associates) were able to confirm inde-

pendently a subset of the results with the data of Loch and colleagues⁵² by using an ANN to distinguish between Gleason grade 3 and 4 lesions. This model correctly identified 82% of grade 3 lesions and 67% of grade 4 lesions.

Crawford and colleagues from the Institute for Clinical Research in Washington, DC (ANNs in CaP Project) are developing software to analyze and interpret TRUS images through pattern recognition software, ANNs, and multivariate analysis. This technology will be developed to identify areas of TRUS images that are indicative of malignant lesions and to distinguish such areas from areas that show healthy tissue. TRUS images are to be gathered prior to RP at three study centers and correlated with the whole-mount prostate specimens. Using digital image processing software, TRUS image cross sections will be matched with their corresponding whole-mount sections, and areas of interest will be marked on the TRUS images. Image filters and pattern recognition software will be used to extract features/signatures from the TRUS images that will then be used to train an ANN. Software is also being developed to construct three-dimensional models from the TRUS cross sections. If these systems can identify malignant lesions within the prostate, a logical next step will be to develop a system to allow real-time direct biopsies of the prostate. This technology may also be applicable to other modalities such as magnetic resonance imaging (MRI) and CDI and may be used in future studies.^{53,54}

The development of technologies to identify malignant lesions in TRUS images using a system of image filters, traditional statistical methods, and ANNs may have a positive impact on prostate cancer patients by improving the accuracy

of needle biopsies. Furthermore, the ability to store three-dimensional images of the prostate may aid patients and their physicians in the watchful waiting strategy.

Repeat Needle Biopsy of the Prostate

Even though serum PSA is the most useful tumor marker for the diagnosis of patients with prostate cancer, this

[Software] will be developed to identify areas of TRUS images that are indicative of malignant lesions and to distinguish such areas from areas that show healthy tissue.

marker is still hampered by a lack of specificity. Until the enhanced ultrasound techniques such as contrast and ANNs become commonplace, many patients will require repeat prostate biopsy.

Approximately two thirds of men undergoing prostate needle biopsy have benign histology. However, a 20% to 40% incidence of positive repeat biopsy in men with elevated PSA who had an initial negative biopsy has been demonstrated.⁵⁵ The presence of prostatic intraepithelial neoplasia or atypia on initial prosta-

9333 men in whom serial PSA measurements were taken, 25% of men with a PSA > 4.0 ng/mL undergoing a second, third, or fourth biopsy following an initial or repeated negative biopsy were found to have prostate cancer.

PSA exists in different molecular forms in the systemic circulation as free PSA, PSA bound to α_1 -anti-chymotrypsin, and PSA bound to α_2 -macroglobulin. The proportion of

PSA α_1 -anti-chymotrypsin was greater in patients with prostate cancer than in those with benign prostatic hyperplasia.^{57,58} Subsequent studies were able to demonstrate some pattern of enhanced specificity without significantly sacrificing sensitivity. A recent multi-institutional trial revealed that patients with a free-to-total PSA ratio of $\leq 10\%$ have a >50% probability of cancer.⁵⁹

One approach toward identifying patients who could benefit from repeat biopsy is the use of the free-to-total PSA ratio as a predictor of

Numerous studies have demonstrated that a significant number of men with an initial negative prostate needle biopsy but persistently elevated serum PSA level will have prostatic malignancy on subsequent biopsy.

tic biopsy are predictive factors of high risk of invasive carcinoma and constitute an indication for repeat biopsy. Numerous studies have demonstrated that a significant number of men with an initial negative prostate needle biopsy but persistently elevated serum PSA level will have prostatic malignancy on subsequent biopsy. In a screening report by Catalona and colleagues⁵⁶ involving

prostate cancer. Brawer and colleagues^{60,61} have studied archival specimens to determine the ability of the free-to-total ratio to predict subsequent prostate cancer. The median Hybritech free-to-total PSA ratio was significantly lower in patients with positive repeat prostate needle biopsy compared with those who had a negative biopsy (14.9% vs 19.4%, $P = .05$). Total PSA as well as the percent

Table 4
Current Indications for Repeat Biopsy

- PSA > 10.0 ng/mL
- cPSA > 7.5 ng/mL
- cPSA/TPSA > 75%
- Strongly suspicious DRE
- Prostate volume > 60 cc

Dianon free-to-Hybritech total PSA ratio were not significantly different between the two groups of men.

For total PSA in the range of 2 to 15 ng/mL, the Hybritech free-to-total PSA ratio appeared to aid in the prediction of cancer on repeat biopsy. A free-to-total PSA ratio in the range of 2 to 15 ng/mL as determined by Hybritech assay appeared to aid in the prediction of cancer on repeat biopsy when biopsy was previously negative. Although this study is limited by the relatively small sample size, the data suggest a potential use of the free-to-total PSA ratio to indicate a higher likelihood for the presence of missed clinically significant carcinoma; hence repeat prostate biopsy would be recommended in clinical practice and other settings. It should be noted that serial free-to-

total PSA ratio measurements can be different among different assays; this should be considered when interpreting serial assays from different labs.^{60,61}

To define the role of repeat needle biopsy further, Brawer and colleagues⁶² presented data on 100 sextant prostate needle biopsies without a diagnosis of malignancy, which were repeated. Carcinoma was detected in 20 repeat biopsies (20%). Stratification based on initial biopsy result revealed carcinoma in 10 of 69 cases (14.5%) without prostatic intraepithelial neoplasia or atypia, 5 of 17 (29.4%) with atypia, 5 of 5 (100%) with grade II or III prostatic intraepithelial neoplasia, and 0 of 9 with grade I prostatic intraepithelial neoplasia. PSA and PSA velocity did not provide statistically significant stratification, perhaps because of the wide variance in these parameters and the small sample size.

These results suggest that patients with a diagnosis of glandular atypia, or grade II or III prostatic intraepithelial neoplasia on initial biopsy, are at high risk for invasive carcinoma and should undergo repeat prostate needle biopsy.⁶² A rapidly increasing serum PSA level or grossly abnormal digital rectal examination may also indicate carcinoma not discovered on initial biopsy (see Table 4).⁶³

Conclusions

Advances continue in the area of early diagnosis of prostate cancer. With the advent of PSA-based screening in the early 1990s, we are witnessing a slow and steady decline in the death rates from prostate cancer. The new tools and approaches reviewed here will continue to fuel improvements in the outcome of patients with prostate cancer. ■

References

1. Eeles RA, the UK Familial Prostate Study Coordinating Group, the CRC/BPG UK Familial Prostate Cancer Study Collaborators. Genetic predisposition to prostate cancer. *Prostate Cancer Prostatic Dis.* 1999;2:9-15.
2. Ostrander EA, Stanford JL. Genetics of prostate cancer: too many loci, too few genes. *Am J Hum Genet.* 2000;67:1367-1375.
3. Eeles RA, Cannon-Albright LA. Familial prostate cancer and its management. In: Eeles RA, Ponder BAJ, Easton DF, Horwich A, eds. *Genetic Predisposition to Cancer.* London: Chapman & Hall; 1996:320-332.
4. Cannon LA, Bishop DT, Skolnick M, et al. Genetic epidemiology of prostate cancer in the Utah Mormon Genealogy. *Cancer Surg.* 1982;1:47-69.
5. Lichtenstein P, Holm NV, Verkasalo PK, et al. Environmental and heritable factors in the causation of cancer—analyses of cohorts of twins from Sweden, Denmark, and Finland. *N Engl J Med.* 2000;343:78-85.
6. Carter BS, Beaty TH, Steinberg GD, et al. Mendelian inheritance of familial prostate cancer. *Proc Natl Acad Sci USA.* 1992;89:3367-3371.
7. Steinberg GD, Carter BS, Beaty TH, et al. Family history and the risk of prostate cancer. *Prostate.* 1990;17:337-347.
8. Bratt O, Borg A, Kristofferson U, et al. CAG repeat length in the androgen receptor gene is related to age at diagnosis of prostate cancer and response to endocrine therapy, but not to prostate cancer risk. *Br J Cancer.* 1999;81:672-676.

Main Points

- Every attempt should be made to detect prostate cancer early, for successful treatment.
- The identification of genetic abnormalities in prostate cancer will allow screening for early disease; some possible gene loci have already been determined.
- Prostate cancer risks include race, family history, and perhaps certain serum markers such as insulin-like growth factor I.
- Transrectal ultrasound (TRUS)-guided biopsy has become the standard for diagnosis.
- Artificial neural networks (ANNs), pattern recognition, and certain contrast agents, all in combination with TRUS, are being explored in order to improve diagnostic accuracy.
- New criteria for follow-up biopsy after an initial negative biopsy are being developed. These include the free-to-total PSA ratio as well as any abnormalities seen on the initial biopsy, such as atypia or high-grade prostatic intraepithelial neoplasia.

9. Giovannucci E, Stampfer MJ, Krithivas K, et al. The CAG repeat within the androgen receptor gene and its relationship to prostate cancer [published erratum appears in *Proc Natl Acad Sci USA*. 1997;94:8272]. *Proc Natl Acad Sci USA*. 1997;94:3320-3323.
10. Hakimi JM, Schoenberg MP, Rondonelli RH, et al. Androgen receptor variants with short glutamine or glycine repeats may identify unique subpopulations of men with prostate cancer. *Clin Cancer Res*. 1997;3:1599-608.
11. Smith JR, Freije D, Carpten JD, et al. Major susceptibility locus for prostate cancer on chromosome 1 suggested by a genome-wide search. *Science*. 1996;274:1371-1374.
12. Berthon P, Valeri A, Cohen-Akenine A, et al. Predisposing gene for early-onset prostate cancer, localized on chromosome 1q42.2-43. *Am J Hum Genet*. 1998;62:1416-1424.
13. Xu J, Meyers D, Freije D, et al. Evidence for a prostate cancer susceptibility locus on the X chromosome. *Nat Genet*. 1998;20:175-179.
14. Gibbs M, Stanford JL, McIndoe RA, et al. Evidence for a rare prostate cancer-susceptibility locus at chromosome 1p36. *Am J Hum Genet*. 1999;64:776-787.
15. Berry R, Schroeder JJ, French AJ, et al. Evidence for a prostate cancer-susceptibility locus on chromosome 20. *Am J Hum Genet*. 2000;67:82-91.
16. Neuhausen SL, Farnham JM, Kort E, et al. Prostate cancer susceptibility locus HPC1 in Utah high-risk pedigrees. *Hum Mol Genet*. 1999;8:2437-2442.
17. Berry R, Schaid DJ, Smith JR, et al. Linkage analyses at the chromosome 1 loci 1q24-25 (HPC1), 1q42.2-43 (PCAP), and 1p36 (CAPB) in families with hereditary prostate cancer. *Am J Hum Genet*. 2000;66:539-546.
18. Xu J, International Consortium for Prostate Cancer Genetics. Combined analysis of hereditary prostate cancer linkage to 1q24-25: results from 772 hereditary prostate cancer families from the International Consortium for Prostate Cancer Genetics [published erratum appears in *Am J Hum Genet*. 2000;67:541-542]. *Am J Hum Genet*. 2000;66:945-957.
19. Tavtigian SV, Simard J, Teng DH, Abtin V, et al. A candidate prostate cancer susceptibility gene at chromosome 17p. *Nat Genet*. 2001;27:172-180.
20. Struwing JP, Hartge P, Wacholder S, et al. The risk of cancer associated with specific mutations of BRCA1 and BRCA2 among Ashkenazi Jews. *N Engl J Med*. 1997;336:1401-1408.
21. The Breast Cancer Linkage Consortium. Cancer risks in BRCA2 mutation carriers. *J Natl Cancer Inst*. 1999;91:1310-1316.
22. Morissette J, Durocher F, Leblanc JF, et al. Genetic linkage mapping of the human steroid 5 alpha-reductase type 2 gene (SRD5A2) close to D2S352 on chromosome region 2p23→p22. *Cytogenet Cell Genet*. 1996;73:304-307.
23. Hankinson SE, Willett WC, Colditz GA, et al. Circulating concentrations of insulin-like growth factor-1 and risk of breast cancer. *Lancet*. 1998;351:1393-1396.
24. Ma J, Pollack MN, Giovannucci E, et al. Prospective study of colorectal cancer risk in men and plasma levels of insulin-like growth factor (IGF)-1 and IGF-binding protein-3. *J Natl Cancer Inst*. 1999;91:620-625.
25. Peehl DM, Cohen P, Rosenfeld RG. The insulin-like growth factor system in the prostate. *World J Urol*. 1995;13:306-311.
26. Pietrowski Z, Mulholland SG, Gomella LG, et al. Inhibition of growth of prostatic cancer cell lines by peptide analogues of insulin-like growth factor 1. *Cancer Res*. 1992;53:1102-1106.
27. Wolk A, Mantzoros CS, Andersson SO, et al. Insulin-like growth factor 1 and prostate cancer risk: a population-based, case control study. *J Natl Cancer Inst*. 1998;90:911-915.
28. Montzoros CS, Tzonou A, Signorello LB, et al. Insulin-like growth factor 1 in relation to prostate cancer and benign prostatic hyperplasia. *Br J Cancer*. 1997;76:1115-1118.
29. Chan JM, Stampfer MJ, Giovannucci E, et al. Plasma insulin-like growth factor-I and prostate cancer risk: a prospective study [see comments]. *Science*. 1998;279:563-566.
30. Kanety H, Madjar Y, Dagan Y, et al. serum insulin-like growth factor-binding protein-2 (IGFBP-2) is increased and IGFBP-3 is decreased in patients with prostate cancer: correlation with serum prostate specific antigen. *J Clin Endocrinol Metab*. 1992;77:229-233.
31. Cohen P, Peehl DM, Stamey TA, et al. Elevated levels of insulin-like growth factor-binding protein-2 in the serum of prostate cancer patients. *J Clin Endocrinol Metab*. 1993;76:1031-1035.
32. Ho PJ, Baxter RC. Insulin-like growth factor-binding protein-2 in patients with prostate carcinoma and benign prostatic hyperplasia. *Clin Endocrinol*. 1997;46:333-342.
33. Schaefer C, Friedman GD. IGF-1 and prostate cancer. *Science*. 1998;282:199a.
34. Sell C, Baserga R, Rubin R. Insulin-like growth factor I (IGF-I) and the IGF-I receptor prevent etoposide-induced apoptosis. *Cancer Res*. 1995;55:303-306.
35. Shariat SF, Bergamaschi F, Adler HL, et al. Correlation of preoperative plasma IGF-I levels with pathologic parameters and progression in patients undergoing radical prostatectomy. *Urology*. 2000;56:423-429.
36. Baffa R, Reiss K, El-Gabry EA, et al. Low serum insulin-like growth factor 1 (IGF-1): a significant association with prostate cancer. *Tech Urol*. 2000;6:236-239.
37. Baserga R, Hongo A, Rubini M, et al. The IGF-I receptor in cell growth, transformation and apoptosis. *Biochim Biophys Acta*. 1997;1332:F105-126.
38. Ismail M, Gomella LG, Alexander AA. Color Doppler sonography of the prostate *Tech Urol*. 1997;3:140-146.
39. Aarnink RG, Beerlage HP, De La Rosette JJ, et al. Transrectal ultrasound of the prostate: innovations and future applications. *J Urol*. 1998;159:1568-1579.
40. Halpern EJ, Strup SE. Using gray-scale and color and power Doppler sonography to detect prostatic cancer. *AJR Am J Roentgenol*. 2000;174:623-627.
41. Presti JC Jr, Chang JJ, Bhargava V, et al. The optimal systematic prostate biopsy scheme should include 8 rather than 6 biopsies: results of a prospective clinical trial [see comments]. *J Urol*. 2000;163:163-166; discussion 166-167.
42. Forsberg F, Merton DA, Liu JB, et al. Clinical applications of ultrasound contrast agents. *Ultrasonics*. 1998;36:695-701.
43. Ragde H, Kenny GM, Murphy GP, et al. Transrectal ultrasound microbubble contrast angiography of the prostate. *Prostate*. 1997;32:279-283.
44. Watanabe M. [Color Doppler enhancement with contrast agents for the detection of prostatic cancer.] *Nippon Rinsho*. 1998;56:1040-1044.
45. Bodgers HA, Sedelaar JP, Beerlage HP, et al. Contrast-enhanced three-dimensional power Doppler angiography of the human prostate: correlation with biopsy outcome. *Urology*. 1999;54:97-104.
46. Halpern EJ, Verkh L, Forsberg F, et al. Initial clinical experience with contrast-enhanced sonography of the prostate. *AJR Am J Roentgenol*. 2000;174:1575-1580.
47. Halpern EJ, Rosenberg M, Gomella LG. Prostate cancer: contrast-enhanced US for detection. *Radiology*. 2001;219:219-225.
48. Dayhoff J. *Neural Network Architectures: An Introduction*. Boston: International Thomson Computer Press; 1996:70-74.
49. Prater JS, Richard WD. Segmenting ultrasound images of the prostate using neural networks. *Ultrasonic Imaging*. 1992;14:159-185.
50. Lizzi FL, Astor M, Feleppa EJ, et al. Statistical framework for ultrasonic spectral parameter imaging. *Ultrasound Med Biol*. 1997;23:1371-1382.
51. Noritomi T, Machi J, Feleppa EJ, et al. In vitro investigation of lymph node metastasis of colorectal cancer using ultrasonic spectral parameters. *Ultrasound Med Biol*. 1998;24:235-243.
52. Loch T, Leuschner I, Genberg C, et al. Artificial neural network analysis (ANNA) of prostatic transrectal ultrasound. *Prostate*. 1999;39:198-204.
53. Ismail M, Petersen RO, Alexander AA, et al. Color Doppler imaging in predicting the biologic behavior of prostate cancer: correlation with disease-free survival. *Urology*. 1997;50:906-912.
54. Miller SM, Ackermann R. Color Doppler sonography of the prostate. *Urol Int*. 1996;57:158-164.
55. Keetch DW, Catalona WJ, Smith DS. Serial prostatic biopsies in men with persistently elevated serum prostate specific antigen values. *J Urol*. 1994;151:1571-1574.
56. Catalona WJ, Smith DS, Ratliff TL, Basler JW. Detection of organ-confined prostate cancer is increased through prostate specific antigen based screening. *JAMA*. 1993;270:948-954.
57. Christenson A, Laurel CB, Lilja H. Enzymatic activity of prostate-specific antigen and its reactions with extracellular serine proteinase inhibitors. *Eur J Biochem*. 1990;194:755-763.
58. Stenman UH, Leinonen J, Alfthan H, et al. A complex between prostate-specific antigen and alpha-1antichymotrypsin is the major form of prostate specific antigen in serum of patients with prostatic cancer: assay of the complex improves clinical sensitivity for cancer. *Cancer Res*. 1991;51:222-226.
59. Catalona WJ, Partin AW, Slawin KM, et al. A multicenter clinical trial evaluation of free PSA in the differentiation of prostate cancer from benign disease [abstract]. *J Urol*. 1997;157:111.
60. Brawer MK, Bankson DD, Haver VM, Petteway JC. Comparison of three commercial PSA assays: results of restandardization of the Ciba Corning method. *Prostate*. 1997;30:269-273.
61. Nixon RG, Petteway JC, Meyer GE, Brawer MK. Comparison of three investigative assays for the free form of prostate-specific antigen [abstract 999]. *J Urol*. 1997;157:255.
62. Brawer MK, Bigler SA, Sohlberg OE, et al. Significance of prostatic intraepithelial neoplasia on prostate needle biopsy. *Urology*. 1991;38:103-107.
63. Carter HB, Pearson JD, Metter EJ, et al. Longitudinal evaluation of prostate-specific antigen levels in men with and without prostate disease. *JAMA*. 1992;267:2215-2220.